

## COURSE HANDOUT

Course Code	ACSC13
Course Name	Design and Analysis of Algorithms
Class / Semester	IV SEM
Section	A-SECTION
Name of the Department	CSE-CYBER SECURITY
Employee ID	IARE11023
Employee Name	Dr K RAJENDRA PRASAD
Topic Covered	Asymptotic Notations
Course Outcome/s	Find the time complexity of an algorithm in terms of different asymptotic notations.
Handout Number	12
Date	10 April, 2023

### Content about topic covered: Asymptotic Notations

#### Asymptotic Notations

**Big O notation:** The function  $f(n) = O(g(n))$  iff (if and only if) there exist positive constants  $c$  and  $n_0$  such that  $f(n) \leq c * g(n)$  for all  $n$ ,  $n \geq n_0$ .

The statement  $f(n) = O(g(n))$  states only that  $g(n)$  is an **upper bound** on the value of  $f(n)$  for all  $n$ ,  $n \geq n_0$ .

Eg:

- |                          |                         |                    |
|--------------------------|-------------------------|--------------------|
| 1. $3n+2 = O(n)$         | $3n+2 \leq 4n$          | $\forall n \geq 2$ |
| 2. $100n + 6 = O(n)$     | $100n + 6 \leq 101n$    | $\forall n \geq 6$ |
| 3. $10n^2+4n+2 = O(n^2)$ | $10n^2+4n+2 \leq 11n^2$ | $\forall n \geq 5$ |
| 4. $6*2^n+n^2 = O(2^n)$  | $6*2^n+n^2 \leq 7*2^n$  | $\forall n \geq 4$ |

Note:  $O(1) \rightarrow$  Constant line

$O(n) \rightarrow$  Linear

$O(n^2) \rightarrow$  Quadratic

$O(n^3) \rightarrow$  Cubic

$O(2^n) \rightarrow$  Exponential




Fig. 1 Big Oh Notation

**Omega notation ( $\Omega$ ):** The function  $f(n) = \Omega(g(n))$  iff (if and only if) there exist positive constants  $c$  and  $n_0$  such that  $f(n) \geq c \cdot g(n)$  for all  $n$ ,  $n \geq n_0$ .

The statement  $f(n) = \Omega(g(n))$  states only that  $g(n)$  is a **lower bound** on the value of  $f(n)$  for all  $n$ ,  $n \geq n_0$ .

Eg:

- |                          |                         |                    |
|--------------------------|-------------------------|--------------------|
| 1. $3n+2 = O(n)$         | $3n+2 \geq 3n$          | $\forall n \geq 1$ |
| 2. $100n+6 = O(n)$       | $100n+6 \geq 100n$      | $\forall n \geq 1$ |
| 3. $10n^2+4n+2 = O(n^2)$ | $10n^2+4n+2 \geq 10n^2$ | $\forall n \geq 1$ |




Fig. 2 Omega Notation




Fig. 3 Theta Notation

**Theta notation ( $\theta$ ):** The function  $f(n) = \theta(g(n))$  iff (if and only if) there exist positive constants  $c_1$ ,  $c_2$  and  $n_0$  such that  $c_1 \cdot g(n) \leq f(n) \leq c_2 \cdot g(n)$  for all  $n$ ,  $n \geq n_0$ .

The statement  $f(n) = \theta(g(n))$  states only that  $g(n)$  is an **both an upper bound and lower bound** on the value of  $f(n)$  for all  $n$ ,  $n \geq n_0$ .

Eg:  $3n + 2 = \theta(n)$        $3n \leq 3n+2 \leq 4n \quad \forall n \geq 1$

**Little Oh notation:** The function  $f(n) = o(g(n))$  iff

$$\lim_{n \rightarrow \infty} \left( \frac{f(n)}{g(n)} \right) = 0$$

Eg:  $3n+2 = o(n^2)$

$$\lim_{n \rightarrow \infty} \left( \frac{3n+2}{n^2} \right)$$

$\rightarrow$

$$\lim_{n \rightarrow \infty} \left( \frac{3}{n} + \frac{2}{n^2} \right) = 0$$